

AD-A042 888

TEXAS UNIV AT AUSTIN DEPT OF ELECTRICAL ENGINEERING  
ASYMPTOTIC PROPERTIES OF CLUSTERING ALGORITHMS, (U)  
JUN 77 L P DEVROYE, T J WAGNER

F/G 12/1

AF-AFOSR-2371-72

UNCLASSIFIED

AFOSR-TR-77-0905

NL

1 of 1  
ADA042888




END

DATE  
FILMED

9 -77

DDC



The U. S. Gov't is authorized to reproduce and sell this report. Permission for further reproduction by others must be obtained from the copyright owner.

Reprinted from the Proceedings of the IEEE Computer Society Conference on Pattern Recognition and Image Processing, Troy, NY, June 6-8, 1977.

ASYMPTOTIC PROPERTIES OF CLUSTERING ALGORITHMS

AFOSR-TR- 77-0905

L.P. Devroye and T.J. Wagner\*  
Department of Electrical Engineering  
The University of Texas at Austin  
Austin, Texas 78712

Abstract

Suppose a sample of size  $n$  is observed from the  $d$ -dimensional density  $f$ . Conditions are given which insure that a single-linkage clustering algorithm can asymptotically find the decomposition of the support of  $f$  into connected closed sets.

Clustering is the process of grouping similar objects. For our purposes the objects to be grouped can be thought of as a set of  $d$ -dimensional vectors and a clustering algorithm can be thought of as any scheme for partitioning this set into subsets called clusters. Our paper analyzes the asymptotic performance of clustering algorithms for a simple probabilistic model with the result that versions of a single-linkage clustering algorithm are shown to be asymptotically effective. Excellent summaries of previous work in clustering are contained in Hartigan and Dorofeyuk<sup>1,2</sup>, while a more technical and thorough description of our results may be found in Devroye and Wagner<sup>3</sup>.

Let  $X$  be a random vector with values in  $\mathbb{R}^d$  and a probability density

$$f = \sum_{i=1}^M \pi_i f_i \quad (1)$$

where  $\pi_i > 0$ ,  $1 \leq i \leq M$ ,  $\sum_{i=1}^M \pi_i = 1$  and  $f_1, \dots, f_M$  are probability densities. If  $f_i$  has support  $C_i$ ,  $1 \leq i \leq M$ , then we assume that

- (a)  $C_i$  is connected,  $1 \leq i \leq M$ ,
- (b)  $C_1, \dots, C_M$  are disjoint, and
- (c)  $C_i$  is bounded,  $1 \leq i \leq M$ .

The supports  $C_1, \dots, C_M$  may be thought of as the clusters chosen by nature. In particular, if independent observations are made on (1) then  $C_1, \dots, C_M$  determine a natural partition of these observations. However, suppose that the statistician assumes only that (1) and (2) hold for some  $M$ ,  $\pi_1, \dots, \pi_M$  and  $f_1, \dots, f_M$ , and, in place of specific knowledge of  $f$ , has a sample size  $n$  from (1), say  $X_1, \dots, X_n$ . The question that concerns us here is how the statistician can asymptotically obtain the same grouping of observations on (1) as he would if he knew  $C_1, \dots, C_M$ .

From the sample  $X_1, \dots, X_n$ , the statistician will, for his clustering algorithm, construct a partition  $A_1, \dots, A_L$  of  $\mathbb{R}^d$ . Future observations, that is, observations from (1) which are independent of those in his sample, will then be grouped together if they fall in the same set  $A_\ell$ . For this reason, we shall

also refer to the sets  $A_1, \dots, A_L$  as clusters. In the vast clustering literature, concentration is focused on grouping the sample  $X_1, \dots, X_n$  and the sets of the partition of  $X_1, \dots, X_n$  determined by  $A_1, \dots, A_L$  are usually referred to as clusters. Concentrating on partitioning  $X_1, \dots, X_n$  seems warranted, for example, in clustering problems arising in paleontology studies where new observations are not expected. However, in medical situations, such as trying to cluster the types of shock for emergency care purposes, the statistician is interested in the performance of the algorithm on future observations. Our model is directed toward this type of situation.

Referring to Figure 1 there are three natural clusters but the algorithm with the sample  $X_1, \dots, X_n$  has yielded four clusters in (a), three in (b) and two in (c). How does one measure the performance of the algorithm on future observations? Agreeing that what we call each cluster  $C_i$  is unimportant as long as we give one unique label to each  $C_i$  we see that the probability of misclassification becomes

$$L_n = \min_g \sum_{i=1}^M \pi_i \int_{A_g(i)} f_i(x) dx \quad (3)$$

where the minimum is taken over all one-to-one functions  $g: \{1, \dots, M\} \rightarrow \{1, \dots, \text{Max}(M, L)\}$  and, if  $M > L$ , we put  $A_{L+1} = \dots = A_M = \emptyset$ . In particular, if  $C_i$  is contained in some  $A_j$  and each  $A_j$  contains at most one  $C_i$  then  $L_n = 0$ . It should be stressed that  $L_n$  is a random variable which depends on  $X_1, \dots, X_n$  and whose value is just the frequency of observations misclassified when a large number of new observations are classified with the partition  $A_1, \dots, A_L$ .

Our interest here is finding what properties are necessary for clustering algorithms to insure that  $L_n \rightarrow 0$  with probability one. The following clustering algorithm, a version of the familiar single-linkage algorithms, has this property with some slight additional assumptions on  $f$ . More extensive results for other algorithms and assumptions may be found in Devroye and Wagner<sup>3</sup>.

If  $r > 0$  connect the two points  $X_i, X_j$  if  $d(X_i, X_j) < r$ ,  $1 \leq i, j \leq n$ . Call two points  $X_k, X_\ell$  connected if there exists a sequence  $Y_0, \dots, Y_m$  from  $\{X_1, \dots, X_n\}$  with  $Y_0 = X_k$ ,  $Y_m = X_\ell$ , and  $Y_{i-1}, Y_i$ , connected,  $1 \leq i \leq m$ . The set  $\{X_1, \dots, X_n\}$  is then partitioned into connected subsets  $K_1, \dots, K_L$ . A partition  $A_1, \dots, A_L$  of  $\mathbb{R}^d$  is obtained from  $K_1, \dots, K_L$  by putting the point  $x \in \mathbb{R}^d$  into  $A_j$  if the closest point to  $x$  from  $X_1, \dots, X_n$  is in  $K_j$  (ties are broken

\*Supported in part by AFOSR Grant 72-2371.

(See 1473)

ADA 042888

AD NU.  
DDC FILE COPY

arbitrarily).

**Theorem.** If  $r = r_n$  satisfies

$$\begin{aligned} (i) \quad nr_n^d / \log n \rightarrow \infty \\ (ii) \quad r_n \rightarrow 0 \end{aligned} \quad (4)$$

and if, for some  $a, b > 0$ ,

$$\inf_{x \in UC_i} \int_{S(x, \rho)} f(x) dx \geq a \rho^d, \quad 0 \leq \rho \leq b, \quad (5)$$

where  $S(x, \rho)$  is the sphere centered at  $x$  with radius  $\rho$ , then

$$L_n \rightarrow 0 \text{ w.p.1.}$$

**Proof.** We recall that the support  $C$  of a density  $f$  is the smallest closed set with the property that

$$\int_C f(x) dx = 1. \text{ In particular, the } C_i \text{ are closed sets.}$$

Because the  $C_i$  are bounded,

$$\inf_{x \in C_i, y \in C_j} \|x - y\| \geq \delta > 0$$

whenever  $i \neq j$ . We assume that  $n$  is so large that  $r_n < \delta$  (use (4(ii))). Suppose that  $X_1, \dots, X_n$  is such that every sphere  $S(x, r_n/3)$  contains at least one of

$$\text{the } X_i \text{ for } x \in \bigcup_{i=1}^M C_i.$$

If we can show that

- (i) whenever  $C_i \cap A_j \neq \emptyset$  and  $X_k \in C_i$ , then  $X_k \in A_j$ , and
- (ii) whenever  $C_i \cap A_j \neq \emptyset$  and  $X_k \notin C_i$ , then  $X_k \notin A_j$ ,

then we know that  $M = L$  and  $\bigcup_{i=1}^M C_i = \bigcup_{i=1}^M (C_i \cap A_{g(i)})$ , for some one-to-one mapping  $g: \{1, \dots, M\} \rightarrow \{1, \dots, M\}$ , which in turn implies that

$$0 \leq L_n \leq \sum_{i=1}^M \pi_i \int_{A_{g(i)}} f_i(x) dx = 0$$

and

$$P\{L_n > 0\} \leq P\{\inf_{x \in C} \mu_n(S(x, r_n/3)) = 0\} \quad (6)$$

where  $\mu_n$  is the empirical measure for  $X_1, \dots, X_n$ .

Let us now prove (i) and (ii). Property (ii) is trivial since  $r_n < \delta$  and  $X_j \in \bigcup_{i=1}^M C_i$  for all  $j$  with probability one. For property (i), we need only show that for any  $x$  in  $C_i$ , and any  $X_j \in C_i$ , there exists a sequence  $Y_1, \dots, Y_\ell$  from  $X_1, \dots, X_n$  with  $Y_1 = x^{(1)}$ ,

$Y_\ell = X_j$ ,  $\|Y_{k+1} - Y_k\| \leq r_n$ ,  $1 \leq k < \ell$ , where  $x^{(1)}$  is the nearest neighbor to  $x$  among  $X_1, \dots, X_n$ .

Since  $S(x, r_n/3)$  contains one  $X_k$ , and since  $r_n < \delta$ , we know that  $x^{(1)}$  belongs to  $C_i$  as well, no matter what  $x$  is picked in  $C_i$ . By the connectedness of  $C_i$ , we can find  $\{x_1, \dots, x_\ell\} \subseteq C_i$  with  $x_1 = x^{(1)}$ ,  $x_\ell = X_j$ , and  $\|x_{k+1} - x_k\| < r_n/3$ ,  $1 \leq k < \ell$ . Thus, since every  $S(x_i, r_n/3)$  contains one of the  $X_k$ 's, we know that there are  $Y_k \in S(x_k, r_n/3)$ ,  $1 \leq k \leq \ell$ ,  $Y_1 = x^{(1)}$ ,  $Y_\ell = X_j$ . Also,  $\|Y_{k+1} - Y_k\| \leq \|Y_{k+1} - x_{k+1}\| + \|x_{k+1} - x_k\| + \|x_k - Y_k\| < r_n$ . This concludes the proof of (i).

As for (6), because the  $C_i$  are bounded, we can find a grid  $\{y_1, \dots, y_N\} \subseteq C$  with the property that for every  $x \in C$  there exists an  $y_i$  with  $\|y_i - x\| \leq r_n/12$ . Such a grid contains at most  $\gamma/r_n^d$  points where  $\gamma > 0$  is a constant depending upon  $d$ ,  $\|\cdot\|$ , and the diameter of  $C$ . If  $r_n/12 < b$ , then

$$\begin{aligned} \inf_i \int_{S(y_i, r_n/6)} f(z) dz &\geq \inf_{x \in C} \int_{S(x, r_n/12)} f(z) dz \\ &\geq a \left(\frac{r_n}{12}\right)^d. \end{aligned}$$

Also, if  $\inf_{x \in C} \mu_n(S(x, r_n/3)) = 0$ , then  $\mu_n(S(y_i, r_n/6)) = 0$  for all  $i$  so that

$$\begin{aligned} P\{L_n > 0\} &\leq P\{\inf_{x \in C} \mu_n(S(x, r_n/3)) = 0\} \\ &\leq \sum_{i=1}^N P\{\mu_n(S(y_i, r_n/6)) = 0\} \\ &\leq \left(\gamma/r_n^d\right) \left(1 - \inf_{x \in \bigcup_{i=1}^M C_i} \int_{S(x, r_n/12)} f(z) dz\right)^n \\ &\leq \left(\gamma/r_n^d\right) \left(1 - r_n^d a/12^d\right)^n \\ &\leq \frac{\gamma}{r_n^d} e^{-\alpha n r_n^d / 12^d}. \end{aligned}$$

By the Borel-Cantelli lemma and (4)(i), we have that  $\sum P\{L_n > 0\} < \infty$ , completing the proof of the Theorem.

Q.E.D.

#### References

- 1 J.A. Hartigan, Clustering Algorithms, John Wiley, N.Y., 1975.
- 2 A.A. Dorofeyuk, "Automatic classification algorithms (review)," Automation and Remote Control, 32, pp. 1928-1958, 1971.
- 3 L.P. Devroye and T.J. Wagner, "Asymptotic Properties of Hierarchical Clustering Algorithms," Technical Report, January 1977, Dept. of Electrical Engineering, University of Texas, Austin, TX.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER <u>18</u> AFOSR-TR-77-0905 ✓	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) <u>6</u> ASYMPTOTIC PROPERTIES OF CLUSTERING ALGORITHMS,	5. TYPE OF REPORT & PERIOD COVERED Interim	
7. AUTHOR(s) <u>10</u> L.P. Devroye and T.J. Wagner	8. CONTRACT OR GRANT NUMBER(s) <u>15</u> ✓ AF- AFOSR 2-2371-72 ✓	
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Texas Department of Electrical Engineering Austin, TX 78712	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F <u>16</u> 2304/A5 <u>17</u> A5	
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research/NM Bolling AFB, Washington, DC 20332	12. REPORT DATE <u>11</u> June 1977 <u>21</u> 4p.	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. NUMBER OF PAGES 2	
	15. SECURITY CLASS. (of this report) UNCLASSIFIED	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) <div style="text-align: right;">D D C REF ID: A61102F AUG 15 1977 A</div>		
18. SUPPLEMENTARY NOTES PROCEEDINGS OF THE IEEE COMPUTER SOCIETY CONFERENCE ON PATTERN RECOGNITION AND IMAGE PROCESSING, Troy, NY, pp. 321-322, June 6-8, 1977.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Clustering; Density Estimation		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Suppose a sample of size $n$ is observed from the $d$ -dimensional density $f$ . Conditions are given which insure that a single-linkage clustering algorithm can asymptotically find the decomposition of the support of $f$ into connected closed sets. <div style="text-align: right;">401997 Incc (over)</div>		



ACCESSION IN	
RTIS	White Section <input checked="" type="checkbox"/>
DRG	Red Section <input type="checkbox"/>
IMPROVED	
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODE	
Dist.	AVAIL. and/or SPECIAL
A	

